

MONTFORT & PRÉAUX

BREEDING REGULATIONS 2019

**PLEASE NOTE : a passport (to include microchip, markings and vaccinations) must accompany the mare on every occasion when walking in.
The following documents must accompany the mare the first time she walks in for covering.**

	Equine Influenza and Equine Herpesvirus	Equine Infectious Anaemia	Equine Viral Arteritis	Contagious Equine Metritis		General Bacterial swab
				CLITORAL	ENDOMETRIAL	
MARES : - BARREN - MAIDEN	<p>COMPULSORY</p> <p>Up to date vaccinations. Last injection done less than a year before the first covering.</p>	<p>RECOMMENDED</p> <p>One negative blood test for imported mares taken after the 1st January 2019.</p>	<p>COMPULSORY</p> <p>Negative result One negative blood test to be taken after the 1st January 2019 and maximum 30 days before 1st service.</p> <p>OR</p> <p>Two Seropositive results The blood tests to be taken after 1st January 2019 a minimum of 14 days apart. The second test has to be taken no longer than 30 days before 1st covering; these should demonstrate a stable or declining titre.</p>	<p>COMPULSORY</p> <p>Negative clitoral swab test for CEMO(Taylorella E.) + Klebsiella P.+ Pseudomonas A.</p> <p>The swab test to be taken after the 1st January 2019</p>	<p>COMPULSORY</p> <p>Negative cervical swab test for CEMO (Taylorella E.) + Klebsiella P.+ Pseudomonas A.</p> <p>The swab test to be taken after the 1st January 2019</p>	<p>RECOMMENDED</p> <p>Negative general bacterial swab test</p> <p>The swab test to be taken after the 1st January 2019</p>
MARE WITH FOAL AT FOOT	<p>COMPULSORY</p> <p>Up to date vaccinations. Last injection done less than a year before the first covering.</p>	<p>RECOMMENDED</p> <p>One negative blood test for imported mares taken after the 1st January 2019.</p>	<p>COMPULSORY</p> <p>Negative result One negative blood test to be taken after the 1st January 2019 and maximum 30 days before 1st service.</p> <p>OR</p> <p>Two Seropositive results The blood tests to be taken after 1st January 2019 a minimum of 14 days apart. The second test has to be taken no longer than 30 days before 1st covering; these should demonstrate a stable or declining titre.</p>	<p>COMPULSORY</p> <p>Negative clitoral swab test for CEMO (Taylorella E.) + Klebsiella P.+ Pseudomonas A.</p> <p>The swab test to be taken after the 1st January 2019, no longer than 30 days before foaling or after the foaling</p>	<p>COMPULSORY</p> <p>Negative cervical swab test for CEMO (Taylorella E.) + Klebsiella P.+ Pseudomonas A.</p> <p>The swab test to be taken after the 1st January 2019 - after the foaling</p>	<p>RECOMMENDED</p> <p>Negative general bacterial swab test</p> <p>The swab test to be taken after the 1st January 2019</p>
HIGH RISK MARE	<p>SEE THE 2018 BREEDING REGULATIONS OF FEDERATION DES ELEVEURS http://www.syndicatdeseleveurs.fr/</p>					